

Automated deep sub-arcsecond wide-field imaging of ELAIS-N1 with LOFAR

Wednesday, 8 May 2024 12:00 (15 minutes)

Recent work by Morabito et al. (2022) and Sweijen et al. (2022) introduced the first steps towards a wide-field imaging pipeline to reduce data from the Low Frequency Array (LOFAR) with all international stations included. Building upon their work, we have improved the pipeline for wide-field imaging by investigating and testing additional pipeline steps to increase the image quality and automation of the pipeline with 32h of LOFAR data of the ELAIS-N1 deep field. This effort led to a wide-field image at a resolution of $0.3''$ and a sensitivity of $17 \mu\text{Jy}/\text{beam}$, marking the deepest wide-field image at this resolution and frequency. In addition, we also generated wide-field images at $0.6''$ and $1.2''$, enabling comparisons of source detections across resolution and sensitivity.

We would like to present this work, as it will bring us closer to the realization of a fully automated LOFAR VLBI pipeline and will allow for the development of surveys similar to the LOFAR Two-metre Sky Survey (LoTSS; Shimwell et al. 2017, 2019, 2022; Williams et al. 2019), improving the ‘standard resolution’ of LOFAR from $6''$ to $0.3''$. This is an important next step to reveal the science hidden among the 90% of radio sources that are currently unresolved in LoTSS.

keywords

radio astronomy, wide-field imaging, VLBI, pipelines, automation, machine learning

In-person or online?

in-person

Career level

ECR

Primary author: DE JONG, Jurjen

Presenter: DE JONG, Jurjen

Session Classification: Techniques